



British Pharmacopoeia Commission Secretariat

MHRA, 10 South Colonnade, Canary Wharf
London, E14 4PU
United Kingdom

British Pharmacopoeia Commission Laboratory

Queen's Road
Teddington, TW11 0LY
United Kingdom

www.pharmacopoeia.com

**BRITISH PHARMACOPOEIA CHEMICAL REFERENCE SUBSTANCE
INFORMATION LEAFLET**

**BUCLIZINE HYDROCHLORIDE IMPURITY STANDARD
CATALOGUE NUMBER 440
CURRENT BATCH: 4516**


Declared Content

No declared content figure is given as the standard is not used for assay purposes.

Use

This British Pharmacopoeia Chemical Reference Substance (BPCRS) is to be used as directed in the monograph(s) of the British Pharmacopoeia and is not intended for any other purpose.

Reference Chromatogram(s)





British Pharmacopoeia Commission Secretariat

MHRA, 10 South Colonnade, Canary Wharf
London, E14 4PU
United Kingdom

British Pharmacopoeia Commission Laboratory

Queen's Road
Teddington, TW11 0LY
United Kingdom

www.pharmacopoeia.com

Related Substances

Column: Nucleosil C₁₈, 10 µm (25 cm x 4.0 mm)

Mobile Phase A: 0.01 M sodium heptanesulfonate in a mixture of 55 volumes of water and 45 volumes of acetonitrile adjusted to pH 4.0 with orthophosphoric acid.

Mobile Phase B: 0.01 M sodium heptanesulfonate in a mixture of 20 volumes of water and 80 volumes of acetonitrile adjusted to pH 4.0 with orthophosphoric acid.

Gradient Programme:	Time (minutes)	Mobile Phase A (%v/v)	Mobile Phase B (%v/v)
	0	100	0
	30	0	100
	40	0	100
	40.5	100	0
	50	100	100

Detection Wavelength: 230 nm

Additional Information

For additional information on BPCRS please visit the British Pharmacopoeia website.

<https://www.pharmacopoeia.com>.