



**BRITISH PHARMACOPOEIA CHEMICAL REFERENCE SUBSTANCE
INFORMATION LEAFLET**

**TRANEXAMIC ACID IMPURITY STANDARD
CATALOGUE NUMBER 734
CURRENT BATCH: 3977**


Declared Content

None given.

Use

This British Pharmacopoeia Chemical Reference Substance (BPCRS) is to be used as directed in the monograph(s) of the British Pharmacopoeia and is not intended for any other purpose.

(b) 1.0% w/v of tranexamic acid impurity standard BPCRS



British Pharmacopoeia Commission Secretariat

MHRA, 10 South Colonnade, Canary Wharf
London E14 4PU
United Kingdom



British Pharmacopoeia

British Pharmacopoeia Commission Laboratory

Queen's Road Teddington, TW11 0LY United
Kingdom

www.pharmacopoeia.com

Related substances

Column: Hypersil ODS (250 mm x 4.6 mm, 5 µm)

Mobile Phase: 11.0 g of anhydrous sodium dihydrogen orthophosphate dissolved in 500 mL of water, 5 mL of triethylamine and 1.4 g of sodium dodecyl sulfate added, pH adjusted to 2.5 with 2 M orthophosphoric acid and diluted to 600 mL with water. 400 mL of methanol added and mixed

Flow Rate: 0.9 mL/minute

Detection Wavelength: 220 nm

Elution order: I: Tranexamic acid (9.0 minutes)

II: Tranexamic acid impurity C (10.2 minutes)

III: Tranexamic acid impurity D (10.6 minutes)

IV: Tranexamic acid impurity A (16.8 minutes)

Note: Tranexamic acid impurity B has a relative retention of about 1.4 (about 12.6 minutes) under these conditions (peak not shown)

Additional Information

For additional information on BPCRS please visit the British Pharmacopoeia website.
<https://www.pharmacopoeia.com>