

Status: Effectivity information can only be shown for content published to the website.

Update information can only be shown for content published to the website.

Levofloxacin Eye Drops

Details for the public consultation of this monograph are as follows:

EAG ABS	Antibiotics
Contact Details	amelia.thomson@mhra.gov.uk peter.crowley@mhra.gov.uk
Deadline for Comment	30 th September 2022
Target Publication (subject to change)	BP 2024
Notes:	New monograph If limits are too restrictive, please provide batch/stability data to demonstrate that an increase is required.

Action and use

Fluoroquinolone antibacterial

DEFINITION

Levofloxacin eye drops are a sterile solution of Levofloxacin Hemihydrate in a suitable vehicle.

The eye drops comply with the requirements stated under [Eye preparations](#) and with the following requirements.

Content of Levofloxacin, $C_{18}H_{20}FN_3O_4$

95.0 to 105.0% of the stated amount.

IDENTIFICATION

- A. Shake a volume of the eye drops containing 0.1g Levofloxacin with an equal volume of [dichloromethane](#). Evaporate the bottom layer to dryness. The [infrared absorption spectrum](#) of the residue, [Appendix II A](#), is concordant with the reference spectrum of Ofloxacin (RS XXX).
- B. In the Assay, the retention time of the principal peak in the chromatogram obtained with solution (1) is similar to that of the peak in the chromatogram obtained with solution (2).

TESTS

[Acidity or alkalinity](#)

pH of the infusion, 6.0 to 7.0, [Appendix V L](#).

Related substances

Carry out the method for [liquid chromatography, Appendix III D](#), using the following solutions.

Solution A: 20% v/v [acetonitrile](#).

- (1) Shake a quantity of the eye drops containing 0.1 g of Levofloxacin with 15 mL of solution A. Dilute to produce 20 mL and filter (0.45- μ m pore size is suitable). Dilute 1 volume to 25 volumes with the mobile phase.
- (2) 0.2% w/v of [levofloxacin BPCRS](#) in solution A, dissolved with the aid of ultrasound if necessary. Dilute 1 volume to 10 volumes with the mobile phase.
- (3) 0.0001% w/v of [levofloxacin impurity B BPCRS](#) in mobile phase.

CHROMATOGRAPHIC CONDITIONS

- (a) Use a stainless steel column (25 cm \times 4.6 mm) packed with [octadecylsilyl silica gel for chromatography](#) (5 μ m) (Inertsil ODS is suitable).
- (b) Use isocratic elution and the mobile phase described below.
- (c) Use a flow rate of 0.8 mL per minute.
- (d) Use a column temperature of 45°.
- (e) Use a detection wavelength of 360 nm.
- (f) Inject 25 μ L of each solution.
- (g) Allow the chromatography to proceed for twice the retention time of levofloxacin.

MOBILE PHASE

0.0874% w/v cupric sulfate, 0.0918% w/v L-isoleucine and 0.594% w/v [ammonium acetate](#) in a mixture containing 3 volumes of [methanol](#) and 7 volumes of [water](#).

SYSTEM SUITABILITY

For system suitability, use solution (2):

the [symmetry factor](#) is not more than 1.8.

CALCULATION OF IMPURITIES

For each unspecified impurity, use the concentration of levofloxacin in solution (1).

For the reporting threshold, use the concentration of levofloxacin in solution (1).

Levofloxacin retention time: about [X] minutes.

Relative retention: impurity E (decarboxy), about 0.38; impurity B, about 0.47; impurity 1 (diamine derivative), about 0.52; impurity C (N-oxide), about 0.63; and impurity A (dextro-), about 1.23.

LIMITS

Identify any peaks corresponding to impurities 1 and E in the chromatogram obtained with solution (1), and multiply the areas of these peaks by the corresponding correction factors: impurity 1, 1.2; impurity E, 1.6.

- impurity A: not more than 0.8%;
- unspecified impurities: for each impurity, not more than 0.2%;
- total impurities, excluding impurity A: not more than 1.0%;

— reporting threshold: 0.1%.

ASSAY

Carry out the method for [liquid chromatography, Appendix III D](#), using the following solutions. Prepare solution A as described under Related Substances.

- (1) Shake a quantity of the eye drops containing 0.5 g of Levofloxacin with 75 mL of solution A. Dilute to produce 100 mL and filter (0.45- μ m pore size is suitable). Dilute 1 volume to 25 volumes with the mobile phase.
- (2) 0.2% w/v of [levofloxacin BPCRS](#) in solution A, dissolved with the aid of ultrasound if necessary. Dilute 1 volume to 10 volumes with the mobile phase.

CHROMATOGRAPHIC CONDITIONS

The chromatographic conditions described under Related substances may be used.

DETERMINATION OF CONTENT

Calculate the content of levofloxacin, $C_{18}H_{20}FN_3O_4$, in the eye drops from the chromatograms obtained and using the declared content of $C_{18}H_{20}FN_3O_4$, in [levofloxacin BPCRS](#).

STORAGE


Levofloxacin Eye Drops should be kept in an airtight container and protected from light.

LABELLING

The quantity of active ingredient is stated in terms of the equivalent amount of Levofloxacin.

IMPURITIES

The impurities limited by the requirements of this monograph include impurities A, B, C and E listed under [Levofloxacin Hemihydrate](#) and:

 1. (S)-9-Fluoro-2,3-dihydro-3-methyl-10-[2-(methylamino)ethylamino]-7-oxo-7H-pyrido[1,2,3-de][1,4]benzoxazine-6-carboxylic acid (diamine derivative)

1. (S)-9-Fluoro-2,3-dihydro-3-methyl-10-[2-(methylamino)ethylamino]-7-oxo-7H-pyrido[1,2,3-de][1,4]benzoxazine-6-carboxylic acid (diamine derivative)